

IN THE CLAIMS:


Amend claims 1, 3, 5-7, 10, 11, 18 and 19 as follows:

61 1. (Twice amended) An administration regimen for improved inhibition of gastric acid secretion [characterized by an extended blood plasma profile of an H^+ , K^+ -ATPase inhibitor,] comprising the oral administration of a pharmaceutical formulation comprising a therapeutically effective amount of an [the] H^+ , K^+ -ATPase inhibitor, wherein the administration regimen induces an extended blood plasma profile of the H^+ , K^+ -ATPase inhibitor [having the formula I




wherein


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

R₆' is hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxycarbonyl, oxazolyl, trifluoroalkyl, or adjacent groups

R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and


R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl].

In claim 3, delete "or 2".

B2
5. (Twice amended) The administration regimen according to claim 1, wherein the extended plasma profile is obtained by oral administration of the pharmaceutical formulation which releases the H^+ , K^+ -ATPase inhibitor for absorption with an almost constant rate during an extended time period and the extended plasma profile is maintained for 2-12 hours.


6. (Twice amended) The administration regimen according to any of claims 1-4 [1-5], wherein the extended plasma profile is maintained for 2-12 hours.

7. (Twice amended) An oral pharmaceutical formulation comprising an H^+ , K^+ -ATPase inhibitor and a pharmaceutically acceptable carrier, wherein the formulation induces an extended blood plasma profile of the H^+ , K^+ -ATPase inhibitor [and the H^+ , K^+ -ATPase inhibitor is a compound of the formula I




wherein


Het₁ is




or




B2
CMT
Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

R₆' is selected from the group consisting of hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;

B² conty
R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups

R₆-R₉ form ring structures which may be further substituted;


R₁₀ is hydrogen or forms an alkylene chain together with R₃; and

R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl].

B³
10. (Twice amended) The oral pharmaceutical formulation according to claim 7, wherein the pharmaceutical formulation releases the H⁺, K⁺-ATPase inhibitor for absorption with an almost constant rate during an extended time period and the extended plasma profile is maintained for 2-12 hours.


11. (Twice amended) The oral pharmaceutical formulation according to any of claims 7-9 [7-10], wherein the extended plasma profile is maintained for 2-12 hours.

B⁴ sub B
18. (Amended) An administration regimen for improved inhibition of gastric acid secretion [characterized by an extended blood plasma profile of an H⁺, K⁺-ATPase inhibitor,] comprising the oral administration of a pharmaceutical formulation comprising a therapeutically effective amount of an [the] H⁺, K⁺-ATPase inhibitor, wherein the administration regimen induces an extended blood plasma profile of the H⁺, K⁺-ATPase inhibitor, [having the formula I




wherein


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

B4
cont'd

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

R₆' is hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups

crit
sub
3


R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and

R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl,]

with the proviso that the H⁺, K⁺-ATPase inhibitor is not pantoprazole.


19. (Amended) An oral pharmaceutical formulation comprising an H⁺, K⁺-ATPase inhibitor and a pharmaceutically acceptable carrier, wherein the formulation induces an extended blood plasma profile of the H⁺, K⁺-ATPase inhibitor [and the H⁺, K⁺-ATPase inhibitor is a compound of the formula I




wherein

B4
CMT
CMT
B3


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

B4
cont
can
sub
B3

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

R₆' is selected from the group consisting of hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;

R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups
R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and


R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl],

with the proviso that the H⁺, K⁺-ATPase inhibitor is not pantoprazole.

Add new claims 20-22.


B5'

20. The administration regimen according to claim 1 or 18, wherein the H⁺, K⁺-ATPase inhibitor is a compound of the formula I




wherein


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

BS
cancel
R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

R₆' is hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;


R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups

R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and


R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl.

C
21. The oral pharmaceutical formulation according to claim 7 ~~or 19~~, wherein the H⁺, K⁺-ATPase inhibitor is a compound of the formula I




wherein


Het₁ is




or




Het₂ is



or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;

B5
Chemist
R₆' is hydrogen, halogen, trifluoromethyl, alkyl and alkoxy;


R₆-R₉ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, halogen, halo-alkoxy, alkylcarbonyl, alkoxy carbonyl, oxazolyl, trifluoroalkyl, or adjacent groups

R₆-R₉ form ring structures which may be further substituted;

R₁₀ is hydrogen or forms an alkylene chain together with R₃; and


R₁₁ and R₁₂ are the same or different and selected from the group consisting of hydrogen, halogen or alkyl.

22. The method according to claim 15 or 16, wherein the H⁺, K⁺-ATPase inhibitor is a compound of the formula I




wherein


Het₁ is




or




Het₂ is




or



X =



or



wherein

N in the benzimidazole moiety means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁, R₂ and R₃ are the same or different and selected from the group consisting of hydrogen, alkyl, alkoxy, fluorine-substituted alkoxy, alkylthio, alkoxyalkoxy, dialkylamino, piperidino, morpholino, halogen, phenyl and phenylalkoxy;

R₄ and R₅ are the same or different and selected from the group consisting of hydrogen, alkyl and aralkyl;